

VACCINE INFORMATION STATEMENT

HPV Vaccine Gardasil® (Human Papillomavirus)

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 What is HPV?

Genital **human papillomavirus (HPV)** is the most common sexually transmitted virus in the United States. More than half of sexually active men and women are infected with HPV at some time in their lives.

About 20 million Americans are currently infected, and about 6 million more get infected each year. HPV is usually spread through sexual contact.

Most HPV infections don't cause any symptoms, and go away on their own. But HPV can cause **cervical cancer** in women. Cervical cancer is the 2nd leading cause of cancer deaths among women around the world. In the United States, about 12,000 women get cervical cancer every year and about 4,000 are expected to die from it.

HPV is also associated with several less common cancers, such as vaginal and vulvar cancers in women, and anal and oropharyngeal (back of the throat, including base of tongue and tonsils) cancers in both men and women. HPV can also cause genital warts and warts in the throat.

There is no cure for HPV infection, but some of the problems it causes can be treated.

2 HPV vaccine: Why get vaccinated?

The HPV vaccine you are getting is one of two vaccines that can be given to prevent HPV. It may be given to both males and females.

This vaccine can prevent most cases of cervical cancer in females, if it is given before exposure to the virus. In addition, it can prevent vaginal and vulvar cancer in females, and genital warts and anal cancer in both males and females.

Protection from HPV vaccine is expected to be long-lasting. But vaccination is not a substitute for cervical cancer screening. Women should still get regular Pap tests.

3 Who should get this HPV vaccine and when?

HPV vaccine is given as a 3-dose series

1st Dose	Now
2nd Dose	1 to 2 months after Dose 1
3rd Dose	6 months after Dose 1

Additional (booster) doses are not recommended.

Routine vaccination

- This HPV vaccine is recommended for girls and boys **11 or 12 years of age**. It *may* be given starting at age 9.

Why is HPV vaccine recommended at 11 or 12 years of age?

HPV infection is easily acquired, even with only one sex partner. That is why it is important to get HPV vaccine before any sexual contact takes place. Also, response to the vaccine is better at this age than at older ages.

Catch-up vaccination

This vaccine is recommended for the following people who have not completed the 3-dose series:

- Females 13 through 26 years of age.
- Males 13 through 21 years of age.

This vaccine *may* be given to men 22 through 26 years of age who have not completed the 3-dose series.

It is *recommended* for men through age 26 who have sex with men or whose immune system is weakened because of HIV infection, other illness, or medications.

HPV vaccine may be given at the same time as other vaccines.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

VACCINE INFORMATION STATEMENT

Tdap Vaccine (Tetanus, Diphtheria, and Pertussis)

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Tetanus, diphtheria and pertussis can be very serious diseases, even for adolescents and adults. Tdap vaccine can protect us from these diseases.

TETANUS (Lockjaw) causes painful muscle tightening and stiffness, usually all over the body.

- It can lead to tightening of muscles in the head and neck so you can't open your mouth, swallow, or sometimes even breathe. Tetanus kills about 1 out of 5 people who are infected.

DIPHTHERIA can cause a thick coating to form in the back of the throat.

- It can lead to breathing problems, paralysis, heart failure, and death.

PERTUSSIS (Whooping Cough) causes severe coughing spells, which can cause difficulty breathing, vomiting and disturbed sleep.

- It can also lead to weight loss, incontinence, and rib fractures. Up to 2 in 100 adolescents and 5 in 100 adults with pertussis are hospitalized or have complications, which could include pneumonia or death.

These diseases are caused by bacteria. Diphtheria and pertussis are spread from person to person through coughing or sneezing. Tetanus enters the body through cuts, scratches, or wounds.

Before vaccines, the United States saw as many as 200,000 cases a year of diphtheria and pertussis, and hundreds of cases of tetanus. Since vaccination began, tetanus and diphtheria have dropped by about 99% and pertussis by about 80%.

2 Tdap vaccine

Tdap vaccine can protect adolescents and adults from tetanus, diphtheria, and pertussis. One dose of Tdap is routinely given at age 11 or 12. People who did not get Tdap at that age should get it as soon as possible.

Tdap is especially important for health care professionals and anyone having close contact with a baby younger than 12 months.

Pregnant women should get a dose of Tdap during every pregnancy, to protect the newborn from pertussis. Infants are most at risk for severe, life-threatening complications from pertussis.

A similar vaccine, called Td, protects from tetanus and diphtheria, but not pertussis. A Td booster should be given every 10 years. Tdap may be given as one of these boosters if you have not already gotten a dose. Tdap may also be given after a severe cut or burn to prevent tetanus infection.

Your doctor can give you more information.

Tdap may safely be given at the same time as other vaccines.

3 Some people should not get this vaccine

- If you ever had a life-threatening allergic reaction after a dose of any tetanus, diphtheria, or pertussis containing vaccine, OR if you have a severe allergy to any part of this vaccine, you should not get Tdap. Tell your doctor if you have any severe allergies.
- If you had a coma, or long or multiple seizures within 7 days after a childhood dose of DTP or DTaP, you should not get Tdap, unless a cause other than the vaccine was found. You can still get Td.
- Talk to your doctor if you:
 - have epilepsy or another nervous system problem,
 - had severe pain or swelling after any vaccine containing diphtheria, tetanus or pertussis,
 - ever had Guillain-Barré Syndrome (GBS),
 - aren't feeling well on the day the shot is scheduled.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention